

# Paper Jurnal UNHAS

Kotak Masuk



**Dyah Kusuma** <herningdkusuma@gmail.com>

Sel, 21 Jun  
2016 07.53

kepada mudhimran, saya

Bersama dengan email ini saya kirimkan paper Ibu Bernadeta Subandini  
untuk Jurnal penelitian di UNHAS

Terimakasih untuk kerjasama yang diberikan

Salam,  
Herning Dyah

# TECTONIC INFLUENCE ON CHANGES IN NEOGENE SEDIMENT SUPPLY, WESTERN PART OF NORTH SERAYU BASIN

Bernadeta Subandini Astuti, ST, M. Eng<sup>\*)</sup>  
Herning Dyah Kusuma W, ST, M.Eng<sup>\*)</sup>  
<sup>\*)</sup>Lecturer of Department of Geology  
Sekolah Tinggi Teknologi Nasional Yogyakarta  
[bernadeta.palguno@gmail.com](mailto:bernadeta.palguno@gmail.com)

## ABSTRACT

Western part of North Serayu Basin composed of Neogene sequence; Halang, Pemali and Rambatan Formation. The oil seeps discovery in this area is an indication of the presence an active petroleum system. Active tectonic condition occur coincidence with the time of deposition. The tectonic influence on change in sediment supply will be the subject of this research.

The combinations of subsidence, eustatic and sediment supply are corresponding to the accommodation space, moreover to the type of sediment which is deposited. The quantity and rate of sediment supply to the basin are closely related to the tectonic activity. In the context of Petroleum Potential, study of sediment supply will provide information about the potential source rock, reservoir and caprock that developed in the research area. During the Neogene at least there have been three times of sea level change in the research area which is related to thickening sedimentary sequence during middle of N18 and thinning in N13 to N17 in some place.

## Introduction

The western part of North Serayu Basin generally composed of Neogene sequences of Halang, Pemali and Rambatan Formation. This basin has potential petroleum system with the discovery of oil seepage. Tectonic condition in the research area during the sedimentation process is still active (Astuti, 2012). The combinations of subsidence, eustatic and sediment supply are corresponding to the accommodation space, moreover to the type of sediment which is deposited. The quantity of sediment supply to the basin closely related to the tectonic activity. In the context of Petroleum Potential, study of sediment supply will provide information about the potential source rock, reservoir dan caprock that developed in the research area. During the Neogene at least there have been three times of sea level change in the research area (Astuti, 2015). Furthermore, Astuti

(2012) mention that the sedimentary sequence is thickening in some research area during middle of N18 and thinning in N13 to N17.

Regional tectonic affecting the reseach area is the Oligo-Miocene north-south trending compression (Satyana, 2007). This tectonic led the formation of Java structural pattern (Pulunggono & Mertodjojo, 1994) with east-west trending reverse fault (Satyana, 2007). Based on van Bemelen (1949), phase of geosyncline evolution occured during Early to Middle Miocene period followed by uplift as geantiklin during Middle to Late Miocene in South Serayu. The formation of geantiklin cause the subsidence in North Serayu geosynclines induced gravitational sliding movement from south direction and generate over thrust in the southern area (van Bemmelen, 1949; Satyana & Armandita, 2006 and Satyana, 2007). Sliding movement in the eastern part of the basin precisely in Karangobar area was repeated in the Pleistocene. In this study we try to analyze the control of tectonic for sedimentation in the research area, since the presence of active tectonic during the sedimentation process will affect the quantity and rate of sediment supply.

### **Data and Research Methods**

The data mainly are collected from Western part of North Serayu area. The stratigraphic and structural data are collected from the field work. These existing data's then analyzed to determine the amount of sediment supply during Neogene. The approach method for analysis is using stratigraphic and sedimentation pattern which will be described as dynamic of sedimentation.

### **Geological Review**

The interaction of eustacy, subsidence, sediment supply, basin physiography, and climate largely control basin sedimentation (Vail et al 1990). One or more variables may be dominant. Different combination of each factor will be related to the relative sea level changes, furthermore to the accomodation space. The accomodation space is the potential space available for sediment to accumulate (after allen, 1997). On the shelf or marine environment, accommodation is controlled by relative sea level, while in fluvial environments;

accommodation is controlled by the fluvial equilibrium profile. Stratigraphic record and its pattern can provide an overview how sedimentary rock is accumulated, factor controlling process and illustrate the dynamic sedimentation.

A discussion of the dynamic sedimentation is closely related with discussion of sedimentology. In general, the package of sedimentary rock is called facies and the set of facies as facies association. These facies or facies association is recorded in stratigraphic column. Stratigraphic column will provide an overview of kind of rocks and its composition, texture, structure, fabric, fossil content and change of stratigraphic element condition vertically and laterally (Rahardjo, 2000).

Sedimentary rocks recording the sedimentation processes during the deposition (Tucker, 1991). Sedimentary rocks formed by the physics, chemistry and biological process in particular sedimentary environment. The physical, chemistry and biological aspect is closely related to the provenance (Boggs, 1992) and characterize its depositional environment both in erosion condition, non deposition and deposition. Product of depositional processes in particular sedimentary environment known as sedimentary facies (Selly, 1985). Sedimentary facies record geometry, lithology, sedimentary structure, fossil content and paleocurrent data.

Regionally the research area included in the 1: 100.000 scale geological map of Majenang area (Kastowo, 1975) published by Geological Research and Development Centre. This geological map represent the regional stratigraphic sucession and structure that developed in research area. Discussion issues related to the sediment supply regionally associated with gliding tectonic mechanism.

Gliding tectonic is variation of gravity tectonic. This tectonic controlled by gravitational force mechanism producing folding and faulting in wide and varying complexity (Allaby & Allaby, 1999 in Satyana 2010). Gravity/vertical tectonic generally associated with fold, thrusting up to gravity sliding from the top to flank of vertically rising fault block, structural arches, mantle, diapirs and like phenomena (Mayerhoff & Hull, 1996)

Sedimentary rocks filling the extensional North Serayu Basin in general are the Neogene turbidities rock with flysch character (Suyanto & Sumantri,

1977). These rocks were deposited on the back arc basin setting. The basin formed by subduction roll back mechanism with intensive structure controlled by rapid and active subsidence during Early to Late Miocene (Koesoemadinata & Martodjojo, 1974).

The extensional process occurred older than N19 until early N20 coincide with subduction process (Astuti, 2012). The rocks were deposited in the western part of the basin during the Middle Miocene to Pliocene from older to younger includes Rambatan, Halang, Pemali, Kumbang and Tapak Formation respectively. These formations generally deposited by debris to turbidity flow mechanism.

The tectonic active during Miocene period influence the sediment deposition rate and change of accomodation space (Koesoemadinata & Martodjojo, 1974). The contrast between the rates of change in accommodation and the sedimentation rates in locations placed in the vicinity of the shoreline allows one to understand why the shoreline may shift either landward or seaward during times of relative sea-level change. Accommodation outpacing sedimentation generates transgression or relative sea level rise, whereas an overwhelming sediment supply may result in shoreline regression or relative sea level fall (Catuneau, 2006).




Figure 1. Stratigraphic compilation by several researchers (Astuti, 2012) show the regional stratigraphic and tectonic.

The relative sea level changes correspond to the shifting of depositional environment during sedimentation process. Shifting of depositional environment recording in stratigraphic record. Sedimentary structure and fossil analysis in western part of North Serayu Basin (Astuti, 2015) indicates there are three times sea level changes during Middle Miocene to Pliocene, i.e., sea level rise, fall and rise.

The sea level rise in N13 to Middle N18 represent by fining upward stacking pattern with depositional environment in the lower fan, upper bathyal. The sea level fall take place in Middle N18 to N19 shown by coarsening upward sequence in the lower to upper fan Upper and Lower Bathyal environment, support with the presence of supra fan lobe. The latest sea level rise start in N19-20 which is indicate with fining upward pattern in the tidal zone to the basin plain or in the neritic to lower bathyal. Relative sea level rise in the research area followed with reducing of sediment supply (Astuti, 2015), while sea level fall is related to erosional process in emerge land and increasing of sediment supply as supra fan lobes.

## **Discussion**

Based on stratigraphic measurements in the study area (Figure 2), the rock sequences in the research area were deposited in upper Neogen (N13 to N20) and distinguished into 10 sequences. N13 to N14 are included in the Middle Miocene, N15 to N18 as the upper Miocene, and N19 to N20 included in the Pliocene. The depositional environment of these facies range from upper or inner fan to basin plain, and partly included tidal zone.

The sediment supply analysis, partly applied with secondary data from some previous researchers. Based on regional study, North Serayu Basin during the Middle Miocene to late Miocene was compressed and uplift which is produce geanticlines in South Serayu (van Bemmelen, 1949, Astuti, 2012). Subsidence in western part of North Serayu takes place at N15 and middle N18 (Figure 1). In the eastern part of the basin, gravitational sliding movements occur from south to north as Nappe Worawari (van Bemmelen, 1949, Astuti, 2016). The provenance is a product of volcanic activity from southern area (Astuti, 2016).




Figure 2. Stratigraphic column of research study in Cikaro River (left) Babakan River (right), were deposited in N18-N20.

The stratigraphic measurement show the decreasing of sediment supply in the eastern part of basin, and increasing in the western part. The age of lithology

in the eastern part of research area based on paleontological analysis is N13 to N17, with erosional event at N15. The erosional event is an evidence of basin uplift which resulted in the absence of deposition. During the middle of N18 the sediment supply increasing especially in Cikaro River and Babakan River in conjunction with the basin subsidence (Astuti 2012, 2015). The combination increasing of sediment supply and basin subsidence caused the accommodation space is constant and the relative sea level is stand still.

Increasing of sediment supply in N19 also occur in the eastern part of research area (Rambatan River). This indicates some uplifted area produce amount of sediment and deposited in the basin afterwards. The sediment covered entire the basin. The increasing sediment supply is not equal with the basin subsidence, however based on Astuti (2015) the eustatic is rise. The combination of these factors indicate that accommodation space is constant start in N18-early N19. During N19-N20 the sediment supply is reduced (Astuti, 2012), the basin subsidence is constant but the eustatic sea level is rise (Astuti, 2015).

## **Conclusion**

Sediment supply based on stratigraphic measurement during N13 to N17 period is minimum, that is followed by uplift and subsidence of the basin. The maximum increasing sediment supply in the research area occurs in N18, and thereafter start from N 19 to N20 the sediment supply is decrease. Increasing sediment supply during N18 followed by basin subsidence and eustatic sea level rise. Decreasing sediment supply takeplace during N19- N20, the basin subsidence is constant, in the other hand the eustatic sea level is rise.



## References

- Astuti, 2012, *Stratigrafi dan Sedimentasi Batuan Neogen di Cekungan Serayu Utara, Daerah Kuningan, Jawa Barat – Larangan, Brebes, Jawa Tengah*, Thesis, tidak dipublikasikan.
- Astuti, 2015, *Perubahan Muka Air Laut Di Cekungan Serayu Utara Bagian Barat Selama Miosen Tengah Hingga Pliosen Di Daerah Kuningan Jawa Barat*, Retti
- Astuti, B. S., 2016, *Identifikasi Provenance Selama Miosen Tengah Hingga Pliosen Di Cekungan Serayu Utara Bagian Barat Di Daerah Kuningan Jawa Barat*, Jurnal Technoscientia, Akprind, Yogyakarta
- Boggs, S. Jr., 2006, *Principles of Sedimentology and Stratigraphy*, fourth edition, Upper Saddle River, New Jersey, p. 662.
- Cateuneanu, 2006, *Principles of Sequence Stratigraphy*, Elsevier
- Kastowo, 1975, *Peta Geologi Lembar Majenang, Jawa, P3G*, Bandung
- Koesoemadinata, R. P. dan Martodjojo, S., 1974, *Penelitian Turbidit di Pulau Jawa*, Laporan research no. 1295174, Badan research Institut Teknologi Bandung, 237 hal.
- Posamentier, H. W., and James, D. P., 1993, *An overview of sequence-stratigraphic concepts : uses and abuses*, *The International Association of Sedimentologists*, Blackwell scientific Publications, p. 3-18.
- Pulunggono, A. & Martodjojo, S., 1994, *Geologi Daerah Pegunungan Selatan: Suatu Kontribusi*, pada : Sriyono., Hendrayana, H., Rahardjo, W. dan Wiyono, S. (Eds.), *Perubahan Tektonik Paleogene-Neogene merupakan peristiwa Terpenting di Jawa*. dalam *Geologi dan Geoteknik Pulau Jawa, sejak akhir Mesozoik hingga Kuartar*; Nafiri, Yogyakarta, Jurusan Teknik Geologi Fakultas Teknik , Universitas Gadjah Mada, hal 37-50.
- Rahardjo, W., 1995, *Foraminifera Besar Tersies Indonesia* (merupakan kumpulan artikel yang diterbitkan Wetenschappelijke Mededeelingen Dienst van Den Minjnbouw in Nederlandch-Indie), Teknik Geologi, Fakultas Teknik UGM.
- Satyana, A.H., dan Armandita, C., 2004, *Deepwater plays of Java, Indonesia : Regional evaluation on opportunities and risks*, IPA Annual Convention Proceedings, DFE04-OR-002, p 27.
- Satyana, A. H., 2007, *Central Java, Indonesia – “A Terra Incognita” in Petroleum Exploration : New Considerations on The Tectonic Evolution and Petroleum Implications*, *Proceedings of Indonesian Petroleum Association Annual Convention*, IPA07-G-085, p. 22.
- Satyana, A. H., 2010, *Gravity Tectonics in Indonesia-A Companion to Plate Tectonics : Cases of Isostatic Exhumation and Gravitational Gliding*, Proceedings PIT IAGI Lombok, p. 12.
- Selley, R.C., 1977, *Ancient Sedimentary Environment*, Cornell University Press, New York, p. 317.
- Sujanto, F.X. dan Sumantri, Y.R., 1977, *Preliminary Study on the Tertiary Depositional Patterns of Java*, IPA Annual Convention Proceedings, p. 183-213.
- Tucker, M.E., 1991, *Sedimentary petrology : an introduction to the origin of sedimentary rocks*, Blackwell scientific publications. P. 260.

- Vail, P. R., Audemard, F., Bowman, S. A., Eiser, P. N, Perezcrus, G., 1990, *The Stratigraphic Signatures of Tectonic, Eustasy and Sedimentation*, AAPG International Lecture, Manuscript version, p. 99.
- Van Bemmelen, 1949, *The Geology of Indonesia; vol. I A, General Geology*; Government Printing Office, The Hague, P. 732.

## Fwd: Paper

Kotak Masuk



**Dyah Kusuma**

10.20 (2 jam  
yang lalu)

kepada saya

----- Forwarded message -----

Dari: **Dyah Kusuma** <[herningdkusuma@gmail.com](mailto:herningdkusuma@gmail.com)>

Date: Sel, 21 Jun 2016 pukul 12.33

Subject: Paper

To: <[mudhimran@gmail.com](mailto:mudhimran@gmail.com)>

Saya coba kirim ulang Pak

Terimakasih

# Manuscript acceptance

Kotak Masuk



**IJEScA <ijesca@unhas.ac.id>**

19 Jul  
2016  
19.50

kepada herringdkusuma, saya

Inggris  
Indonesia

[Terjemahkan pesan](#)

[Nonaktifkan untuk: Inggris](#)

Dear Bernadeta Subandini Astuti, ST, M. Eng

Thank you for submitting the manuscript, "TECTONIC INFLUENCE ON CHANGES IN NEOGENE SEDIMENT SUPPLY, WESTERN PART OF NORTH SERAYU BASIN" to

International Journal of Engineering and Science Applications.

The Article is accepted as a part of the IJEScA vol. 3, 1, May 2016.

If you have any questions, please contact me. Thank you for considering this journal as a venue for your work.

Thank you for your participation

A. M. Imran

International Journal of Engineering and Science Applications

---

International Journal of Engineering and Science Applications  
<http://pasca.unhas.ac.id/ojs/index.php/ijesca>