

# SIM-Nanggulan formation as

*by Al Hussein Flowers Rizqi Al Hussein Flowers Rizqi*

---

**Submission date:** 12-Dec-2022 10:46PM (UTC-0500)

**Submission ID:** 1979817120

**File name:** journal\_44-3\_21\_1.pdf (1.15M)

**Word count:** 3985

**Character count:** 22997

Original Article

Nanggulan formation as a roof pendant at the central part of  
Kulon Progo Mountains, Yogyakarta, Indonesia

Winarti<sup>1\*</sup>, Al Hussein Flowers Rizqi<sup>1</sup>, and Emi Sukiyah<sup>2</sup>

<sup>1</sup> Department of Geology, Faculty of Mineral Technology,  
Institut Teknologi Nasional Yogyakarta, Sleman Regency, Yogyakarta, 55281 Indonesia

<sup>2</sup> Department of Geology, Faculty of Geological Engineering,  
Universitas Padjadjaran, Bandung, 40132 Indonesia

Received: 13 July 2021; Revised: 27 December 2021; Accepted: 29 December 2021

Abstract

The Nanggulan Formation sediments form a weakly undulating hill and range in age from the Middle Eocene to the Late Oligocene. In Samigaluh and Sermo areas, a series of sedimentary rocks with small dimensions resembling fragments and forming a steep hilly morphology is exposed. Outcrops are mainly made up of sandstone and claystone interbedded with lignite intercalation. An outcrop in Sermo-2 found andesite intruding claystone. Claystone is found near rock contact, where it is an alteration to metasediment and chlorite is formed. Petrographic data showed the presence of chlorite, pyrite, and kaolinite. Andesite dike is one factor that contributes to the Nanggulan Formation being exposed to the surface. The Nanggulan Formation is a roof pendant above a dike. The implications of the roof pendant are related to the propylitic to argillic of hydrothermal alteration, which produce chlorite, pyrite, illite, and kaolinite minerals.

**Keywords:** roof pendant, metasediment, fragments, dike, alteration

1. Introduction

The Nanggulan Formation, as the oldest formation in the Kulon Progo Mountains, has a minimal spread. This formation is only exposed on the Kulon Progo Mountains eastern side (Widagdo, Pramumijoyo, & Harijoko, 2018). Surface geological studies show that the Nanggulan Formation, which is exposed in the eastern part of the Kulon Progo Mountains, is caused by a thrust fault, sinistral fault, dextral fault, and anticline (Widagdo, Pramumijoyo, & Harijoko, 2020). The existence of a thrust fault on the surface lends support to gravity research on Sentul and Mujil lines (Winarti, Sukiyah, Syafri, & Nur, 2020). The Nanggulan Formation, exposed on the eastern side of the Kulon Progo Mountains, is relatively widespread.

The Nanggulan Formation, which is exposed around Mount Mujil, is caused by a pressure factor from the Old Andesite Formation with brittle and rigid properties, known as a tumor-like bulge (Winarti, Sukiyah, Syafri, & Nur, 2021b). In the eastern Kulon Progo Mountains, Mount Gede found tuff that was exposed after becoming broken through by igneous rock, resulting in the tuff forming a roof pendant above the broken through rock. It is characterized by tuff in igneous rock as xenolith (Bronto, Ratdomopurbo, Asmoro, & Rani, 2014).

The Nanggulan Formation is found on weakly undulated hills, while the Old Andesite Formation is found on steep hills (Winarti, Sukiyah, Syafri, & Nur, 2020). The Nanggulan Formation will generally be at the bottom, while the Old Andesite Formation will be at the top. If the Nanggulan Formation is found above the Old Andesite Formation and has the morphology of steep hills, then it is influenced by factors.

Sedimentary rocks exposed between andesite breccias in the central part of the Kulon Progo Mountains

\*Corresponding author

Email address: winarti.itny@gmail.com

raise two questions: are these fragments of the Nanggulan Formation, and how can these sedimentary rocks be exposed above the Old Andesite Formation? This research aims to reveal the factors that cause sedimentary rock to be constructed in the central Kulon Progo Mountains (Samigaluh and Sermo), the implications for hydrothermal alteration zoning and mineralization processes.

2. Materials and Methods

2.1 Geological setting

The Kulon Progo Mountains, which are part of the micro-continent Gondwana, is assumed to be a structural boundary that extends northeast-southwest to Mount Muria (Smyth, Hamilton, Hall, & Kinny, 2007). These boundaries have an irregular shape and are associated with complex tectonics. In contrast, the Kulon Progo Mountains have many structural lineaments which trend northwest-southeast, west-east, north-south, and northeast-southwest (Smyth, Hamilton, Hall, & Kinny, 2007). Some of the lineament of structure develops into faults, and so do some of the triggering intrusions (Widagdo, Paramumijono, Harijoko, & Setiawan, 2016).

The Kulon Progo Mountains comprise of the Nanggulan Formation, the Old Andesite Formation, the

Jonggrangan Formation, the Sentolo Formation, and volcanic deposits (Widagdo, Paramumijono, Harijoko, 2018). The Nanggulan Formation comprises clastic sedimentary rocks such as sandstone, claystone, quartz sandstone with lignite, and calcareous claystone intercalation. The rocks have plastic properties (Winarti, Sukiy, Syafri, & Nur, 2020). The Nanggulan Formation ages from the Middle Eocene to Early Upper Eocene (Saputra & Akmaluddin, 2015), and the Middle Eocene to Late Oligocene (Coxall *et al.*, 2021).

Overlaying the Nanggulan Formation is the Old Andesite Formation, which is comprised of volcanic rocks (andesitic intrusion and andesitic lava, with andesite breccia) (Winarti, Sukiyah, Syafri, & Nur, 2021a). The age of the Old Andesite Formation is Late Oligocene to Middle Miocene (Soeria-Atmadja *et al.*, 1994), and early Oligocene (Coxall *et al.*, 2021).

Sedimentary rock outcrops in Samigaluh and Sermo are found above andesite dike and occupy a steep hill. If plotted on a regional geological map, it is relatively located in the middle of the Kulon Progo Mountains (Figure 1). The sedimentary rock outcrop is surrounded by volcanic rocks known as the Kebobotak Formation (Tomk) and the Andesite (a) (Rahardjo, Sukandarrumidi, & Rosidi, 2012) or the Old Andesite Formation (Bemmelen, 1949).




Figure 1. Samigaluh (WSG) and Sermo (SRM-2) are located in the central Kulon Progo Mountains, which are part of the Kebobotak Formation (Tomk) and Andesite (a).

## 2.2 Intrusion and alteration zone

When magma mineralizes before reaching the surface, it causes intrusion (Martí, Groppelli, & Silveira, 2018). Plutonic rocks that break through the rock series above have a non-conformity relationship (Martí, Groppelli, & Silveira, 2018). The intrusion that breaks through rock walls can impact hydrothermal alteration of the porphyry system (Sukisman, Yoyok, Mulyaningsih, & Hidayah, 2021). Complex geological structures can also trigger hydrothermal alteration of porphyry types (Adriansyah, Zakaria, Muslim, & Hirnawan, 2018).

Rudnik Mts. is a volcano-intrusive complex with shallow intrusions associated with mineralization (Cvetković, Šarić, Pécskay, & Gerdes, 2016). The intrusion contains many xenoliths of schists, gneiss and amphibole fragments as a side effect of contact metamorphism. Metamorphic rocks of the Pinoh Formation in the Schwane Mountains become large-scale roof pendants due to tonalite intrusions, characterized by the presence of metamorphic xenoliths in tonalite (Nurdini, Febyani, & Budiman, 2012). Roof pendants are commonly found as small fragments and crushed rock due to intrusion (Stevens & Greene, 1999).

The roof pendant at La Medrelena mine is 300 meters long and 50 meters wide (Nokleberg, 1981). According to theory, a roof pendant is a rock exposed on the surface due to a relatively shallow intrusion (volcanic plutonic), and rocks that have undergone metamorphism such as metasediment quartzite and schists (Dunning, Walstrom, & Lechner, 2018). The Andesite intrusion in the Kulon Progo Mountains has weathered and parts have experienced hydrothermal alteration in propylitic and argillic phases (Phyu, Hendrayana, Indrawan, & Kamai, 2021). The intrusion at Mt. Gede that forms a roof pendant is characterized by silica and argillic tuff as a mineral altered from pumice lapilli, coarse tuff, and fine tuff (Bronto, Ratdomo, 1990, Asmoro, & Rani, 2014).

Based on the mineral assemblage produced, the alteration is classified as propylitic type (chlorite, epidote, carbonate), argillic type (smectite, montmorillonite, illite-smectite, kaolinite), low temperature advanced argillic type (kaolinite, alunite), advanced temperature argillic type high (pyrophyllite, diasporic, andalusite), potassic type (adularia, biotite, quartz), phylic type (quartz, sericite, pyrite), sericytic type (sericite, quartz, muscovite), and silicified type (quartz) (Guilbert, & Park, 1986).

According to Seager *et al.* (1982 vide Evans, 1987), the propylitic zone develops in the outer and peripheral alteration zones and is characterized by mineral assemblages of epidote, calcite, and chlorite. When  $H^+$  and  $CO_2$  are added, the primary mafic minerals (biotite and hornblende) are altered to chlorite and carbonate. This zone is also dominated by metal sulfides such as pyrite. The argillic zone is distinguished by mineral assemblages of clay, quartz, and carbonate. Pyrite can be found in trace amounts. Argillic is found in the hydrothermal system's outer layer.

## 2.3 Methods

Methods used in this research involved field observations, micropaleontology analysis, petrography, and X-Ray Diffraction (XRD) analysis. Micropaleontology analysis was carried out using a binocular microscope, the

Olympus CX-31 P, with a 10x magnification. This micropaleontology analysis aimed was to identify planktonic and benthic foraminifera fossils, which are used to determine the relative age of rocks and their depositional environment.

Petrographic analysis was carried out using a polarized microscope Olympus with a magnification of 40x. Petrography analysis aimed to identify the presence of mineralization that could be used as an indicator of a metamorphic process. Samigaluh (WSG) rock samples are sandstone and claystone, whereas Sermo (SRM) rock samples are sandstone, claystone, and andesite.

The goal of XRD analysis is to determine the types of minerals that are not identified by petrography analysis. An XRD analysis was performed on altered rocks to identify the type of mineral that resulted from the alteration. The alteration zone is determined by minerals formed as a result of the alteration.

## 3. Results and Discussion

### 3.1 Observations in the field

Based on field observations, the sedimentary rock outcrop in Samigaluh has a height of 6–7 meters and a width of 20 meters. Overall, outcrops are composed of interbedded sandstone and claystone, with lignite intercalation and brittle rock properties (Figure 2). Sandstone layer thickness ranges from 8 to 50 cm, claystone layers range from 7 to 30 cm, and lignite layer thickness is 8 cm, with strike and dip of rock bedding N026E/05. The slope of the small rock layers (relatively horizontal) indicates whether the rock is still in normal condition or disturbed by tectonics. At the bottom, the sedimentary rock is surrounded by andesite breccia, a volcanic product rock.

The sedimentary rock outcrop at Sermo-2 is also composed of interbedded sandstone and claystone with coal intercalation, with strike and dip of rock bedding N210E/30. The rock can be found above the andesite dike (Figure 3A). A chloritization zone forms between the two rocks' contacts (Figure 3B).

The sandstone is brownish-gray, layered, and composed primarily of quartz. Claystone is characterized by its incredibly dark color, complex properties, shale structure, extensive fracture, and the rock has been altered into metasediment. A normal fault cuts through sedimentary rocks at Sermo (SRM-2), and sedimentary rocks overlap above andesite dike in a stratigraphic position. Metasediment occurs when an andesite dike comes into contact with claystone, causing claystone to change into metasediment and chlorite present (Figure 3B). An andesite dike produced the head.

The dike in the field is characterized by igneous rock breaking through claystone, with the majority of the igneous rock weathered. Dike has a massive, interlocking texture, is holocrystalline, and is composed of mafic minerals (mainly pyroxene and biotite), with feldspar partially altered into clay minerals.

### 3.2 Petrographic analysis

Petrographic analysis of Sermo-2 (SRM-2) sandstone reveals that the grain shape is anhedral-subhedral, grain supported, and well sorted. The rocks are composed of






Figure 2. Sandstone and claystone interbedded with lignite intercalation




Figure 3. A. An Andesite Dike Old Andesite Formation intrudes sedimentary rock (Nanggulan Formation) at Sermo (SRM-2). B. Chlorite as a characteristic of the metamorphism process caused by the andesite dike at Sermo (SRM-2).

minerals such as feldspar, quartz, carbon, pyrite, chert, and rock fragments such as metaquartz and granitic fragments. Kaolinite exists to serve as a block of cement and is usually formed through intermediate alteration processes (Figure 4A). The presence of chlorite, pyrite, and kaolinite minerals in sandstones can indicate hydrothermal alteration.

Petrographic analysis of igneous rocks at Sermo-2 (SRM-2) revealed that the rocks have a porphyritic texture with moderate aphanitic to faneric granularity, a degree of crystallization including holocrystalline, relatively subhedral-euhedral mineral form, and phenocryst embedded in the base mass of a finer mineral. The rocks have a unique pilotactic texture, which is indicative of shallow intrusions.

The rocks contain minerals such as pyroxene, plagioclase, sanidine, chlorite, and opaque minerals with a plagioclase microlite base mass (Figure 4B). The presence of biotite, sanidine, and chlorite minerals as evidence of hydrothermal alteration occurred in that location.




Figure 4. Petrographic analysis results at Sermo-2 (cross-polarized light position). A. Sandstone indicated the existence of pyrite and kaolinite, which indicate hydrothermal alteration. B. Chlorite minerals indicate andesite dike as evidence of hydrothermal alteration.

The results of claystone petrographic analysis at the Sermo-1 (SRM-1) location show that grain forms are anhedral-subhedral, matrix-supported, and composed of feldspar, carbon, quartz, and pyrite (Figure 5). In this section, microfractures can be seen. The presence of pyrite indicates that the claystone has been hydrothermally altered too. The sedimentary rocks in Samigaluh and Sermo, composed of sandstone and claystone interbedded with lignite intercalation, are well layered, and their brittle rock properties are similar to those of the Nanggulan Formation, based on physical characteristics of the rock and petrographic analysis.

### 3.3 X-Ray diffraction analysis

Rock samples for XRD analysis were collected from an altered andesite dike at Sermo (SRM-3). The XRD analysis results (Figure 6), show that the constituent minerals are mostly illite (90.4%) and kaolinite (9.6%).

### 3.4 Micropaleontology Analysis

The analysis of the relative age of sandstone and claystone from outcrop at Samigaluh (WSG) discovered fossils of planktonic foraminifera *Globigerapsis kugleri*, *Globigerinatheka barri*, *Globorotalia centralis*, and *Globigerina mexicana* (Table 1). Based on the fossil planktonic foraminifera content, the rock age range is Middle Eocene (P14).



Table 3. Benthic foraminifera fossils found in Samigaluh show that the rock was deposited in an outer Neritic.

No	Species	Neritic			Bathyal		Abysal	
		Inner (0-20) m	Middle (20-100) m	Outer (100-200) m	Upper (200-500) m	Middle (500-1000) m	Outer (900-2000) m	(2000-5000) m
1	<i>Cibicides parki</i>							
2	<i>Textularia zeaguta</i>							
3	<i>Rotaliatina sulcigerina</i>							

Table 4. Benthic foraminifera fossils found in Sermo show that the rock was deposited in an inner Neritic.

No	Species	Neritic			Bathyal		Abysal	
		Inner (0-20) m	Middle (20-100) m	Outer (100-200) m	Upper (200-500) m	Middle (500-1000) m	Outer (900-2000) m	(2000-5000) m
1	<i>Virgulina pontomi</i>							
2	<i>Cibicides sp</i>							

Based on relative ages of rocks, sedimentary rocks in Samigaluh and Sermo that are similar to the Nanggulan Formation are Eocene in age. So, based on outcrop data and the age of the rock, it can be determined whether sedimentary rock outcrops in Samigaluh and Sermo are part of the Nanggulan Formation. However, the rocks exposed in two locations are smaller than the Nanggulan Formation in general, and are in the form of fragments. Furthermore, based on field data and petrographic analysis, it is believed that the Nanggulan Formation in Samigaluh and Sermo has undergone local metamorphism caused by andesite dike. These metamorphisms produce metasediments and minerals such as chlorite, illite, sanidine, pyrite, and kaolinite.

The Nanggulan Formation, which rode above the Old Andesite Formation and is exposed as a result of a dike, has become a roof pendant in the central part of the Kulon Progo Mountains (Figure 7). So, if the Nanggulan Formation, which is exposed in the central part of the Kulon Progo Mountains, is controlled by volcanism in the form of dike, this becomes evidence.




Figure 7. Model of the Nanggulan Formation roof pendant on the central side of Kulon Progo

#### 4. Conclusions

The Nanggulan Formation is the oldest formation in Kulon Progo Mountains. This formation is mostly exposed on the eastern side of the Kulon Progo Mountains, and tectonics is the main controlling factor. Sedimentary rocks exposed at Samigaluh and Sermo (in the central Kulon Progo Mountains) and forming fragments between volcanic rocks are confirmed to be part of the Nanggulan Formation.

A shallow andesite dike intrudes the sedimentary rocks, forming a chloritization zone at the contact between the two rocks. Furthermore, the heat generated by the andesite dike, claystone, and sandstone changes into metasediment and the form of alteration minerals (chlorite, pyrite, illite, and kaolinite).

The presence of chlorite mineral, as determined by outcrop and petrographic data, characterizes the type of propylitic alteration that occurs in the system's outer zone. The presence of metal sulfides such as pyrite also helps support this zone. Illite and kaolinite identified from XRD analysis are most commonly formed in the low-temperature advanced argillic type alteration, the same as the hydrothermal alteration system's outer zone.

The Nanggulan Formation, which was uplifted to the surface due to andesite dike intrusion, is also characterized by undisturbed rock layers (relatively horizontal). The sediment rocks (Nanggulan Formation) function as a roof pendant to the above dike (Old Andesite Formation). To determine the depth and shape of the dike beneath the surface, a geophysical survey (gravity or magnetic) is required.

#### Acknowledgements

The author would like to thank the team that assisted with data collection in the field.

## References

- Adriansyah, Y., Zakaria, Z., Muslim, D., & Hirnawan, F. (2018). Determining of geotechnical domain based on joint density and fault orientation at batu hijau mine, west sumbawa-Indonesia. *International Journal of Engineering, Transactions A: Basics*, 31(4), 679–683. doi:10.5829/ije.2018.31.04a.21
- Bemmelen, R. W. V. (1949). *Geology of Indonesia Vol-IA general*. Leiden, the Netherlands: The Hague: Martinus Nijhoff.
- Bronto, S., Ratdomopurbo, A., Asmoro, P., & Rani, M. A. (2014). Gigantic landslides of Merapi Volcano, Yogyakarta-Central Java. *Journal of Geology and Mineral Resources*, 15(4), 165-183. doi:10.33332/jgsm.geologi.v15i4.
- Coxall, H. K., Jones, T. D., Jones, A. P., Lunt, P., MacMillan, I., Mariyani, G. I., N. . . Pearson, P. N. (2021). The eocene–oligocene transition in Nanggulan, Java: lithostratigraphy, biostratigraphy and foraminiferal stable isotope. *Journal of the Geological Society*. doi:10.1144/jgs2021-006
- Cvetković, V., Šarić, K., Pécskay, Z., & Gerdes, A. (2016). The rudnik mts. volcano-intrusive complex (central serbia): An example of how magmatism controls metallogeny. *Geologia Croatica*, 69(1), 89–99. doi:10.4154/gc.2016.08
- Dunning, G. E., Walstrom, R. E., & Lechner, W. (2018). Barium silicate mineralogy of the western margin, north American continent, part 1: Geology, origin, paragenesis and mineral distribution from Baja California Norte, Mexico, Western Canada and Alaska, USA. *Baymin Journal*, 19(5), 1–70. Retrieved from <https://www.researchgate.net/publication/328232378>.
- Evans, A. M. (1993). *Ore geology and industrial minerals, An introduction* (3<sup>rd</sup> ed.). Hoboken, NJ: Blackwell Science.
- Guilbert, J. M., & Park, C. F. Jr. (1986). *The geology of ore deposits*. New York, NY: W. H. Freeman and Company.
- Martí, J., GropPELLI, G., & Silveira, A. B. D. (2018). Volcanic stratigraphy: A review. *Journal of Volcanology and Geothermal Research*, 357, 68–91. doi:10.1016/j.jvolgeores.2018.04.006
- Nokleberg, W. J. (1981) Stratigraphy and structure of the strawberry mine roof pendant central Sierra Nevada, California. Washington, DC: United States Government.
- Nurdini, W. P., Febyani, S., & Budiman, A. R. (2012). Optimization of non-volcanic geothermal potential in West Kalimantan using hot dry rock (HDR) method. *Proceeding of National Earth Seminar 5<sup>th</sup>*, E11-1 – E11-17.
- Phyu, H. T., Hendrayana, H., Indrawan, I. G. B., & Kamai, T. (2021). Simulation of Kalirejo Road side slope based on altered andesite characters, Kulon Progo Regency, Indonesia. *Journal of Applied Geology*, 5(2), 101. doi:10.22146/jag.56916
- Rahardjo, W., Sukandarrumidi & Rosidi, H. M. D. (2012). *Geological map Yogyakarta sheet, Java 1: 100.000*. Bandung, Indonesia: Geological Research and Development Centre.
- Saputra, R., & Akmaluddin. (2015). Biostratigrafi nannofossil gampingan Formasi Nanggulan bagian bawah berdasarkan batuan inti dari Kec. Girimulyo dan Kec. Nanggulan, Kab. Kulon Progo, D.I. Yogyakarta. *Proceeding of National Earth Seminar 8<sup>th</sup>*, 400-412. Retrieved from <https://repository.ugm.ac.id/id/eprint/135460>
- Smyth, H. R., Hamilton, P. J., Hall, R., & Kinny, P. D. (2007). The deep crust beneath island arcs: Inherited zircons reveal a Gondwana continental fragment beneath East Java, Indonesia. *Earth and Planetary Science Letters*, 258, 269–282. doi:10.1016/j.epsl.2007.03.044
- Soeria-Atmadja, R., Maury, R. C., Bellon, H., Pringgoprawiro, H., Polve, M., & Priadi, B. (1994). Tertiary magmatic belts in Java. *Journal of Southeast Asian Earth Sciences*, 9(1–2), 13–27. doi:10.1016/0743-9547(94)90062-0
- Stevens, C. H., & Greene, D. C. (1999). Stratigraphy, depositional history, and tectonic evolution of paleozoic continental-margin rocks in roof pendants of the eastern Sierra Nevada, California. *Bulletin of the Geological Society of America*, 111(6), 919–933. doi:10.1130/0016-7606(1999)111<0919:SD HATE>2.3.CO;2
- Sukisman, R. S. Yoyok, Mulyaningsih, S., Hidayah, A. R. (2021). Hydrothermal alteration and ore metal mineralisation at Temon, Pacitan, East Java, Indonesia. *Journal of Geoscience, Engineering, Environment, and Technology*, 6(1), 24–33. doi:10.25299/jgeet.2021.6.1.6368.
- Watt, S. F. L. (2019). The evolution of volcanic systems following sector collapse. *Journal of Volcanology and Geothermal Research*, 384, 280–303. doi:10.1016/j.jvolgeores.2019.05.012
- Widagdo, A., Paramumijono, S., Harijoko, A., & Setiawan, A. (2016). Preliminary study of geological structures control for the distribution of rocks in Kulonprogo Mountain Region Yogyakarta. *Proceedings of National Earth Seminar 9<sup>th</sup>*, 9–20. Retrieved from <https://repository.ugm.ac.id/id/eprint/137861>
- Widagdo, A., Pramumijoyo, S., & Harijoko, A. (2018). The morphotectono-volcanic of Menoreh-Gajah-Ijo volcanic rock in western side of Yogyakarta-Indonesia. *Journal of Geoscience, Engineering, Environment, and Technology*, 3(3), 155. doi:10.24273/jgeet.2018.3.3.1715
- Widagdo, A., Pramumijoyo, S., & Harijoko, A. (2020). Geological structure control toward Naggulan formation presence in the Naggulan district, Kulon Progo Regency, Yogyakarta. *Journal of GEOSAPTA*, 6(2), 97. doi:10.20527/jg.v6i2.8282
- Winarti, Sukiyah, E., Syafri, I., & Nur, A. A. (2020). Springs phenomena as contacts between nanggulan and old andesite formations at eastern west progo dome, Indonesia. *International Journal of Geomate*, 19(74), 167–175. doi:10.21660/2020.74.81033



Winarti, Sukiyah, E., Syafri, I., & Nur, A. A. (2021a). The comparative of morphological and gravity anomaly lineaments in West Progo Mountains, Indonesia. *International Journal on Advanced Science Engineering Information Techonology*, 11(1), 363-368. doi:10.18517/ijaseit.11.1.13076

Winarti, Sukiyah, E., Syafri, I., & Nur, A. A. (2021b). The Nanggulan formation and old andesite formation contact at Mujil Mount, West Progo-Yogyakarta. *KURVATEK*, 6(1), 117-124. doi:10.33579/krvtk.v6i1.2086

# SIM-Nanggulan formation as

## ORIGINALITY REPORT

11%

SIMILARITY INDEX

7%

INTERNET SOURCES

9%

PUBLICATIONS

2%

STUDENT PAPERS

## PRIMARY SOURCES

- 1 [ijog.geologi.esdm.go.id](http://ijog.geologi.esdm.go.id) 1%

Internet Source
- 2 [journal.itny.ac.id](http://journal.itny.ac.id) 1%

Internet Source
- 3 Muhammad Nurjati Setiawan, Jatmika Setiawan, Alim Sugiarto. "Slope stability analysis for development of simple housing in Pandhawa Hill, Jering, Godean, Sleman", AIP Publishing, 2020 1%

Publication
- 4 Yulaikhah, Subagyo Pramumijoyo, Nurrohmat Widjajanti, Asmoro Widagdo. "Optimal design of the Sermo Fault deformation monitoring network using sensitivity criteria based on geological information", Arabian Journal of Geosciences, 2021 1%

Publication
- 5 Asmoro Widagdo, Subagyo Pramumijoyo, Agung Harijoko, Agung Setiyanto. "Fault lineaments control on disaster potentials in 1%

# Kulon Progo Mountain Area-Central Java-Indonesia", MATEC Web of Conferences, 2018

Publication

---

6	Submitted to Universitas Riau Student Paper	1 %
7	repositorio.ucv.edu.pe Internet Source	1 %
8	Nurul Dzakiya, F D Sastrawan, R B Laksana, M M Anan Amara SA. "Identification of lithology properties of groundwater by using resistivity method in Girimulyo, Kulon Progo, Yogyakarta", Journal of Physics: Conference Series, 2019 Publication	1 %
9	www.insightsociety.org Internet Source	1 %
10	gsm.org.my Internet Source	1 %
11	Agung Septana Nugraha, Darsono Darsono, Budi Legowo. "Identification of the distribution of andesite rocks in Kalirejo Village, Kokap District, Kulon Progo Regency, Special Region of Yogyakarta based on geoelectrical tomography data", Journal of Physics: Conference Series, 2019 Publication	<1 %

---

12

Helen K. Coxall, Tom Dunkley Jones, Amy P. Jones, Peter Lunt et al. "The Eocene–Oligocene transition in Nanggulan, Java: lithostratigraphy, biostratigraphy and foraminiferal stable isotopes", Journal of the Geological Society, 2021

Publication

<1 %

13

Muhamad Afdal Ahmad Basri, Shazlyn Milleana Shaharudin, Kismiantini, Mou Leong Tan et al. "Regionalization of Rainfall Regimes Using Hybrid RF-Bs Couple with Multivariate Approaches", ISPRS International Journal of Geo-Information, 2021

Publication

<1 %

14

[searg.rhul.ac.uk](http://searg.rhul.ac.uk)

Internet Source

<1 %

15

A Widagdo, S Pramumijoyo, A Harijoko. "Tectonostratigraphy-volcanic of Gajah-ljo-Menoreh Tertiary volcanic formations in Kulon Progo mountain area, Yogyakarta-Indonesia", IOP Conference Series: Earth and Environmental Science, 2018

Publication

<1 %

16

[www.mdpi.com](http://www.mdpi.com)

Internet Source

<1 %

17

[digitallibrary.usc.edu](http://digitallibrary.usc.edu)

Internet Source

<1 %



18

A Widagdo, S Pramumijoyo, A Harijoko.  
"Morphotectono-volcanic of Tertiary volcanic  
rock in Kulon Progo mountains area,  
Yogyakarta-Indonesia", IOP Conference  
Series: Earth and Environmental Science, 2018  
Publication

---

<1 %

19

Andi Faesal, Muhammad Irman Khalif Ahmad  
Aminuddin, Aji Syailendra Ubaidillah. "Host  
rock petrology, hydrothermal alteration  
characteristics & ore mineralogy of porphyry  
copper-gold deposit, Brambang, Lombok,  
West Nusa Tenggara Indonesia", Materials  
Today: Proceedings, 2022  
Publication

---

<1 %

20

[journal.ugm.ac.id](http://journal.ugm.ac.id)  
Internet Source

---

<1 %

Exclude quotes  On

Exclude matches  Off

Exclude bibliography  On